Geometry A

•

7.6 : Apply Sine and Cosine Ratios Notes

Name: ______ Date: ______

Period:


- I can use sine and cosine ratios to find missing side lengths in right triangles.
- I can apply trigonometric ratios to real-world problems.

In the last section, we looked at the tangent ratio for an acute angle in a right triangle, which involved only the lengths of the two legs of a right triangle. The **sine** and **cosine** ratios are ratios for acute angles in right triangles that involve the length of a ______ and the ______ of the right triangle.

Trigonometric Ratios	
Let $\triangle ABC$ be a right triangle with acute $\angle A$, then the sine of $\angle A$ (abbreviated sinA) is defined as:	5 4
$sinA = \frac{length of leg opposite \angle A}{length of hypotenuse}$	
Let $\triangle ABC$ be a right triangle with acute $\angle A$, then the cosine of $\angle A$ (abbreviated cosA) is defined as:	5 4
$\cos A = \frac{\text{length of leg adjacent to } \angle A}{\text{length of hypotenuse}}$	


Example 1: Find sine ratios

Find sinX and sinY. Write each answer as a fraction in simplest form and as a decimal rounded to four places.




Example 2: Find cosine ratios.


Find cos*P* and cos*Q*. Write each answer as a fraction in simplest form and as a decimal rounded to four places.



Example 3: Use trigonometric ratios to find side lengths


Use a trigonometric ratio to find the value of x in the diagram. Round answer to nearest tenth.





Example 4: Apply trigonometric ratios to real world situations

a) A rope staked 20 feet from the base of a building goes to the roof and forms an angle of 58° with the ground. To the nearest tenth of a foot, how long is the rope?



b) Michael, whose eyes are six feet off the ground, is standing 36 feet away from the base of the building, and he looks up at a 50° angle of elevation to a point on the edge of the building's roof. To the nearest foot, how tall is the building?

