



- I can perform calculations using the midpoint formula.
 - I can calculate the distance between two points.
 - I can apply the midpoint formula to solve a context problem.

Bisector: Line RS bisects \overline{PQ} at point R. Find RQ if $PQ = 4\frac{3}{4}$ inches.



Midpoint Formula: 1 dimension

$$M = \frac{x_1 + x_2}{2}$$

1. Find the midpoint of the line segment.



2. Using Midpoints: In the diagram, M is the midpoint of the segment. Find DE.



Midpoint Formula: 2 dimensions

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

3. Find the midpoint of a line with the given endpoints: A(4, -3) and B(5, 6)

4. Caliyaah is traveling to Peru for her summer vacation. She looks at a map of the path of her flight. Her plane leaves from Georgia, located at $(5, 16)$ on the map's coordinate grid, makes a stop at the halfway point, Panama, and then heads to its destination, Peru, located at $(7, 8)$ on the map's coordinate grid. Find the location of Panama on Caliyaah's map.
5. Find the second endpoint of the line with the given endpoint (P) and midpoint (M).
- a) P(8, 0) and M(6, -5)
 - b) P(7, -17) and M(-2, 3)




- I can perform calculations using the distance formula.
 - I can calculate the distance between two points.
 - I can apply the distance formula to solve a context problem.

Distance Formula: 1 dimension

$$D = |x_2 - x_1|$$

1. Find the length of the line segment.



Distance Formula: 2 dimensions

$$D = \sqrt{x_2 - x_1^2 + y_2 - y_1^2}$$

2. Find the distance between two given points.

a) $A(3, 5)$ and $B(5, 7)$

b) $R(2, 3)$ and $S(4, -1)$

Extension: Is $\overline{AB} \cong \overline{RS}$? Explain ☺

3. In the diagram to the right, is the distance from Joan's home to school the same as the distance from Starbuck's to Joan's home? Explain.

